

74T-E34. RONALD FAGIN, IBM T. J. Watson Research Center, Yorktown Heights, New York 10598.

$L_{\kappa, \lambda}$ -equivalence of ordinals and a compactness result.

Throughout $\kappa \geq \lambda \geq \omega$, κ regular; α, β, γ are ordinals. *Definitions.* 1. Write α uniquely as $\gamma\alpha_1 + \alpha_2$, $\alpha_2 < \gamma$. Then $r_\gamma(\alpha)$ and $c_\gamma(\alpha)$ are defined to be α_2 and cofinality $(\gamma\alpha_1)$, respectively. Further, $\alpha \sim \beta \pmod{\gamma}$ if (1) $r_\gamma(\alpha) = r_\gamma(\beta)$, and (2) $\alpha < \gamma$ iff $\beta < \gamma$. 2. $\mathcal{U} \equiv \mathcal{B}$ means that \mathcal{U} and \mathcal{B} agree on $L_{\kappa, \lambda}$ -sentences.

Theorem. 1. If λ is a successor or ω , $\kappa = \lambda$, then $\langle \alpha; \epsilon \rangle \equiv_{\kappa, \lambda} \langle \beta; \epsilon \rangle$ iff (1) $\alpha \sim \beta \pmod{\kappa^\kappa}$, and (2) $c_{\kappa^\kappa}(\alpha)$ and $c_{\kappa^\kappa}(\beta)$ are equal, or both at least λ . 2. If λ is regular, $\kappa > \lambda$, then $\langle \alpha; \epsilon \rangle \equiv_{\kappa, \lambda} \langle \beta; \epsilon \rangle$ iff (1) $\alpha \sim \beta \pmod{\kappa}$, and (2) $c_\kappa(\alpha)$ and $c_\kappa(\beta)$ are equal, or both at least λ . 3. If λ is weakly inaccessible, $\kappa = \lambda$, then $\langle \alpha; \epsilon \rangle \equiv_{\kappa, \lambda} \langle \beta; \epsilon \rangle$ iff (1) $\alpha \sim \beta \pmod{\kappa}$, and (2) $c_\kappa(\alpha)$ and $c_\kappa(\beta)$ are equal, or both at least λ . 4. If λ is singular, $\kappa = \lambda^+$, then $\langle \alpha; \epsilon \rangle \equiv_{\kappa, \lambda} \langle \beta; \epsilon \rangle$ iff (1) $\alpha \sim \beta \pmod{\kappa^\kappa}$, and (2) $c_{\kappa^\kappa}(\alpha)$ and $c_{\kappa^\kappa}(\beta)$ are equal, or both at least λ^+ . 5. If λ is singular, $\kappa > \lambda^+$, then $\langle \alpha; \epsilon \rangle \equiv_{\kappa, \lambda} \langle \beta; \epsilon \rangle$ iff (1) $\alpha \sim \beta \pmod{\kappa}$, and (2) $c_\kappa(\alpha)$ and $c_\kappa(\beta)$ are equal, or both at least λ^+ . \square The theorem yields a complete characterization of ordinals in $L_{\kappa, \lambda}$ (using only $<$ and $=$), from which follows *Corollary.* Assume $\kappa > \lambda$, or κ is weakly inaccessible; $\kappa = \lambda$. Let Σ be a set of $L_{\kappa, \lambda}$ sentences involving only $<$ and $=$, \exists every subset of Σ having less than κ sentences has a well-ordered model. Then Σ has a well-ordered model. (Received January 7, 1974.)